

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## Nova Aquilæ No. 2.

A single magnitude observation of *Nova Aquilæ*, made December 16th, completes my series for 1905. (See these *Publications*, October and December.) Further measures cannot be made here till the latter part of February.

The estimates were made with the 12-inch, power 155, by the Argelander method, at a large hour-angle  $(5^h.1)$  under poor atmospheric conditions. Four photometric settings on the star f, with the 12-inch, November 25th, gave for its magnitude 13.46.

G. M. T., 1905. Estimates. Nova. Dec.  $16^{\rm d}.60$  e 5-6 x 3 Nova 12 f; d 8 e 8 Nova 12 $^{\rm m}$ .1 The value of a step from this observation is  $0^{\rm m}.085$ . January 25, 1906. James D. Maddrill.

## RAINFALL AT MT. HAMILTON.

Until the 9th of January the season beginning 1905, July, threatened to be a dry one. From January 11th to 19th, however, almost continuous rain brought the season record up from 4.93 inches to 17.23 inches. The heaviest fall

In one hour was 0.77<sup>in</sup>, from 12:40 to 1:40 P. M., January 12th.

In 24 consecutive hours was 3.86in, from 3 P. M., January 11th, to 3 P. M., January 12th.

Of the latter 3.39 inches fell in the last 12 hours. The next day the fall

In 24 consecutive hours was 3.83in, from 6 A. M., January 13th, to 6 A. M., January 14th.

Of this 3.61 inches fell in 12 hours, from 7 A. M. to 7 P. M.

The twenty-five-year summary at the conclusion of this article shows that about 14.5 inches are to be expected by January 19th; hence we are about 2.7 inches ahead of the normal season.

A table of the rainfall by months for the first twelve years, 1880, July – 1892, June, was compiled by Mr. Perrine in 1893 and published in these *Publications* (Vol. V, p. 126). The following table is a continuation to 1905, June; and a summary

in the last two columns based on all the rainfall data for the twenty-five years.

Melted snow is included with rain.

Month.	1892 1893	1893 1894	1894 1895	1895 1896	1 <b>8</b> 96 1 <b>89</b> 7	1897 1898	1898 1899	1899 1900
July	in. 0.00	in. 0.00	in. 0.02	in. O.OI	in. 0.00	in. 0.00	in. 0.00	in. 0.00
August	0.00	0.00	0.00	0.00	0.28	0.00	0.00	0.12
September	0.24	0.48	1.64	0.08	0.47	0.07	0.29	0.00
October	1.38	o.66	2.98	0.78	1.85	1.25	1.33	6.37
November	10.30	4.01	0.84	2.46	5.86	1.51	1.23	4.92
December	5.56	3.58	11.90	3.16	4.91	2.70	2.13	4.16
January	3.29	9.74	10.00	9.54	3.50	2.30	5.63	3.26
February	3.45	10.52	3.08	1.08	7.42	4.16	0.75	1.70
March	8.99	2.54	1.46	3.83	6.45	2.04	11.11	3.37
April	3.61	0.89	2.30	6.7c	0.82	0.84	1.40	4.06
May	0.95	2.78	2.39	2.10	0.28	2.41	1.47	1.35
June	0.16	0.64	0.00	0.02	0.38	0.38	0.39	0.00
Annual	37.93	35.84	36.61	29.76	32.22	17.66	25.73	29.31
Month.	1900	1901 1902	1902 1903	1903 1904	1904 1905	25-year		lean Sun to End of Month.
т 1	in.	in.	in	in.	in.	in.		in.
July	0.01	0.00	0.00	0.00	0.00	0.00		0.00
August	0.02	0.05	0.00	0.00	0.05	0.03		0.03
September	0.08	1.08	0.00	0.00	2.33	0.41		0.44
October	3.48	2.19	2.09	0.47	2.51	1.82		2.26
November	7.76	2.89	3.01	7.69	2.05	3.01		5.27
December	2.21	1.61	3.11	1.39	3.84	6.53		11.80
January	5.76	1.44	8.86	1.98	4.04	4.67		16.47
February	5.92	9.15	2.20	9.53	3.89	4.72		21.19
March	1.98	5.18	9.89	8.06	5.91	5.39		26.58
April	3.33	2.60	1.12	4.38	1.36	3.11		29.69
May	1.07	1.19	0.05	0.45	2.27	1.63		31.32
June	0.02	0.00	0.00	0.03	0.00	0.38		31.70
Annual	31.64	27.38	30.33	33.98	28.25	31.2	 70	

January 22, 1906.

JAMES D. MADDRILL.

## A Double-Star Problem.

Burnham has called 95 Ceti (= Alvan Clark 2) "the most mysterious and strange double star in the heavens. I have tried it," he says, "first and last, perhaps hundreds of times with apertures all the way from 6 to 36 inches without